



REVIEW ARTICLE

# Status of Hepatozoonosis in Wildlife: A Global Overview

Sayed Jafar Hasani<sup>a,b</sup>, Bijan Esmailnejad<sup>c,\*</sup>, Farzin Abbasi<sup>d</sup>, Awat Samiei<sup>e</sup>, Navid Azarm<sup>d</sup>, Mohammad Reza Najafi<sup>c</sup>, Negin Esfandiari<sup>f</sup>, Seyyed Erfan Hasani<sup>g</sup>

<sup>a</sup> Department of Basic Medical Sciences, khoj University of Medical Sciences, khoj, Iran.

<sup>b</sup> Pathogens and Vectors Research Center, Cellular and Molecular Medicine Research Institute, Urmia University of Medical Sciences, Urmia, Iran.

<sup>c</sup> Department of Pathobiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran.

<sup>d</sup> DVM graduate, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran.

<sup>e</sup> Food and Beverages Safety Research Center, Urmia University of Medical Sciences, Urmia, Iran.

<sup>f</sup> Department of Food Hygiene and Quality Control, Division of Epidemiology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran.

<sup>g</sup> DVM student, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran.

## Article info

### Article history:

Received 2025-12-05

Received in revised form

2025-12-12

Accepted 2025-12-13

### Keywords:

Tick-Borne Protozoan

Wildlife

*Hepatozoon* spp.

## Abstract

*Hepatozoon* spp. are parasites that are transmitted by ticks and can cause a range of diseases in both wild and domestic animals, varying in severity. These parasites have the ability to infect a diverse range of species such as birds, reptiles, amphibians, and mammals. They are frequently encountered in tropical and subtropical areas on all continents. Various carnivores, such as foxes (*Vulpes vulpes*), black-backed jackals (*Canis mesomelas*), golden jackals (*Canis aureus*), African wild dogs (*Lycaon pictus*), hyenas (*Crocuta crocuta*), cheetahs (*Acinonyx jubatus*), leopards (*Panthera iridus*), and lions (*Panthera leo*), have been identified as hosts for similar species of *Hepatozoon* based on their physical characteristics. This article aims to provide a comprehensive review of the findings regarding *Hepatozoon* spp. in both free-roaming and captive wild animals from America, Europe, and Asia. The discovery of *Hepatozoon* spp. in different types of mammals, including those that are domesticated and those that are wild, indicates that there is a possibility of diseases spreading between animals that live in the wild and those that are domesticated.

## 1. Introduction

The investigation of *Hepatozoon*, a genus containing over 300 species of apicomplexan parasites, has primarily focused on domestic animals and peri-urban areas, neglecting its occurrence in natural environments and wild hosts, which remains poorly explored (Ferrari *et al.*, 2022). The *Hepatozoon* genus includes a variety of species that can infect a diverse array

of animals, spanning birds, reptiles, amphibians, and mammals. These infections are prevalent in tropical and subtropical regions across all continents (O'Dwyer, 1907). until a decade ago, histology methods were primarily used to study blood parasites, with a few exceptions. Nevertheless, since the mid-2010s, molecular methods have become increasingly popular for detecting and identifying *Hepatozoon* infections. Specifically, researchers have utilized the amplification of fragments

\*Corresponding author: B. Esmailnejad (Associate Professor)

E-mail address: b.esmailnejad@urmia.ac.ir


<http://dx.doi.org/10.22084/avr.2025.32041.1028>

of the small subunit ribosomal RNA (18S rRNA) gene. These advanced techniques have enabled the identification of numerous *Hepatozoon* species in wild mammals throughout Europe, America, and Asia, with a particular emphasis on carnivores. Although the 18S rDNA gene has been commonly used as a molecular marker for phylogenetic analyses, problems in separating closely related species might occur due to its high degree of conservation. All *Hepatozoon* species share a common life cycle, which includes gametogony and sporogony taking place in the definitive host, typically a bloodsucking invertebrate. In the intermediate host, usually a vertebrate, schizogony occurs, followed by the development of gametes (Hamšíková *et al.*, 2016). *Hepatozoon* spp. can be transmitted in multiple ways, but the most common method is through the ingestion of infected vectors or hosts. In mammals, for example, animals can become infected when they consume arthropod ectoparasites like ticks, fleas, and mosquitoes, which carry the parasite and are attached to another individual. This transmission often occurs during grooming behavior (Hamšíková *et al.*, 2016; Johnson *et al.*, 2010; Paperna *et al.*, 2004; Rigó *et al.*, 2016). Furthermore, transmission can happen within predator-prey food webs when predators consume prey that either carry infectious meronts or are infested by infected vectors. Additionally, vertical transplacental transmission has also been observed (Duscher *et al.*, 2015; Murata *et al.*, 1993). *Hepatozoon* spp. have been found in a wide range of habitats across the globe. These habitats include temperate broad-leaved forests, conifer forests, Mediterranean scrublands, subtropical forests, savannahs, and taiga. These findings suggest the presence of intricate transmission cycles that remain largely unexplored (Battisti *et al.*, 2020; Duscher *et al.*, 2014; Laakkonen *et al.*, 2001; Modrý *et al.*, 2021; Perles *et al.*, 2019; Tomé *et al.*, 2014). However; there is still a significant gap in our understanding of the occurrence and frequency of *Hepatozoon* spp. in essential ecosystems. Particularly, limited research has been conducted in exploring their presence across different altitude levels, where the swift habitat changes correspond to a diverse range of hosts that coexist to some extent. Mountainous habitats are highly vulnerable to both abiotic and biotic changes caused by global climate alterations. These changes can lead to the movement of mammal species to higher elevations, which in turn can cause significant alterations within local communities (Gilman *et al.*, 2010; Urban *et al.*, 2016). Consequently, there is a possibility for the expansion of parasites in both their geographic range and the variety of host species they infect (Garcia-Vozmediano *et al.*, 2020; Rizzoli *et al.*, 2019). Monitoring the presence of *Hepatozoon* spp., a parasite with a highly adaptable transmission cycle, is vital in habitats experiencing rapid changes. In rodents, *Hepatozoon* spp. infection can be pathogenic, but typ-

ically presents as a mild and asymptomatic condition (Brumpt *et al.*, 1946; Hoogstraal, 1961; Johnson *et al.*, 2008; Miller, 1908; Simpson *et al.*, 2006). In contrast, carnivorous animals like canids, felids, mustelids, and snakes may experience a weakened immune system due to *Hepatozoon* spp. This makes them more vulnerable to co-infection with other pathogens (Alencar *et al.*, 1997; Cunningham and Yabsley, 2011; Diakou *et al.*, 2020; Zechmeisterová *et al.*, 2021). This article aims to provide a comprehensive review of the findings regarding *Hepatozoon* spp. in both free-roaming and captive wild animals from America, Europe, and Asia. The sampling methods and materials used, as well as the laboratory detection methods employed, will be discussed in detail.

## 2. Reports of *Hepatozoon* Species in Domestic and Wild Canids

**Dogs.** Two species of *Hepatozoon* have been discovered in pet dogs: *Hepatozoon canis*, which has been observed in various regions including Europe, Asia, Africa, South America, and the United States; and *Hepatozoon americanum*, which has only been found in the United States so far. In Brazil, the only species found affecting dogs is *H. canis* (Forlano *et al.*, 2007; Vincent-Johnson *et al.*, 1997; Shkap *et al.*, 1994). *H. canis*, a protozoan belonging to the phylum Apicomplexa, was initially discovered in dogs' blood in India and identified as *Leukocytozoon canis* (Christophers, 1907; James, 1905). *H. canis* like *Theileria* spp. and *Rangelia vitalii* can infects both leukocytes and parenchymal tissues (Fig. 1).



**Fig. 1.** Giemsa-stained blood smear showing gamonts of *Hepatozoon* spp in neutrophils (100x).

Dogs become infected with *H. canis* when they consume ticks that harbor mature oocysts (Baneth *et al.*, 2001). The brown dog tick, known as *Rhipicephalus sanguineus*, plays a vital role as the main host for *H. canis*. Although in Brazil, the main vector of *H. canis* is *Amblyomma ovale*. Dogs and wild canids, on the other hand, serve as intermediate hosts in this process.<sup>3</sup> In

general; there is a significant difference in how common *Hepatozoon* spp. is between red foxes and dogs living in the same areas (Cassini *et al.*, 2009; Conceição-Silva *et al.*, 1988; Gabrielli *et al.*, 2010). Blood PCR analysis revealed the presence of *H. canis* in 21.1% of the 331 dogs surveyed in kennels and shelters located in southern Portugal (Menn *et al.*, 2010). Foxes are considered to be important carriers of *H. canis* for pet dogs. It is interesting to observe that foxes have been present in the world for a much longer time than the domestication of dogs (Lindblad-Toh *et al.*, 2005; Allen *et al.*, 2008). Conducted a study to examine the Hepatozoon species found in the blood of 274 domestic dogs in Oklahoma, United States. Nested PCR and Real-time PCR techniques were employed for the analysis. The findings of the study indicated that *H. americanum* was the most commonly found species in dogs residing in endemic areas, as well as in clinically ill and shelter dogs. Moreover, certain dogs (n=2) that presented with clinical signs of American canine hepatozoonosis or similar organisms referred to Oklahoma State University. These results indicate that *H. americanum*, *H. canis*, and *H. canis* could potentially be responsible for causing diseases in dogs residing in the southern part of the United States (Allen *et al.*, 2008). A study conducted in Brazil in 2016 by de Castro Demoner *et al.* uncovered a significant occurrence of *H. canis* in dogs. By employing both microscopic examination and molecular analysis, they determined that around 66.45% of the dogs were infected with *H. canis* (de Castro Demoner *et al.*, 2016). *H. canis* infection is highly prevalent in dogs, and it is commonly found in areas where the tick vector *R. sanguineus* is present. This encompasses various regions, including Africa, Southern Europe, South America, Asia (including the Middle East), as well as islands in the Pacific and Indian Oceans (Table 1).

Hepatozoonosis is a disease that is commonly found in animals and its prevalence can vary. Previous research has shown different percentages of *H. canis* seroprevalence in dogs. For example, in Portugal, it was reported to be 36%, in Nigeria 17.6%, in India 2.5%, and in Thailand 2.1% (Gevrey, 1993). In order to accurately evaluate the degree and seriousness of *H. canis* infection in dogs, it is crucial to conduct comprehensive and systematic studies. This is because the infection often does not show any clinical signs. Furthermore, the occurrence of *H. canis* infection greatly differs in various regions. For example, 39% of dogs in rural areas of Rio de Janeiro state, Brazil, were found to have circulating *H. canis* gamonts in their blood. In Zaria, Nigeria, this percentage was 22% among dogs, whereas in Malaysia, only 1.2% of dogs were affected (Ezeokoli *et al.*, 1983; O'dwyer *et al.*, 2001; Rajamanickam *et al.*, 1985). The PCR analysis did not thoroughly investigate the similarity among various *H. canis* isolates. A thorough examination of the 18S rRNA of Hepatozoon

isolates from dogs in Japan and Israel showed a significant resemblance, with a similarity of up to 99% to *H. canis*. However, these isolates were distinct from *H. americanum* (Inokuma *et al.*, 2002). The initial clinical case of *H. canis* infection in Bulgaria was documented by Ivanov and Kanakov (Ivanov and Kanakov, 2003). Jittapalapong *et al.* were the first to molecularly confirm the presence of *H. canis* in Thailand (Jittapalapong *et al.*, 2006). Canine hepatozoonosis was initially discovered in 1978 around the Gulf Coast area of Texas, marking the first documented cases in the United States. Since then, the illness has been found in dogs throughout a large part of the southeastern United States. At first, it was believed to be caused by *H. canis*, but the disease showed a much more severe progression compared to cases of canine hepatozoonosis seen in other parts of the world. Significant variations were noted not just in the clinical symptoms and laboratory abnormalities, but also in the pathological characteristics, tissue specificity, parasite structure, and tick vectors. In 1997, this organism was officially acknowledged as a separate species called *H. americanum* (Craig *et al.*, 1978). Javad Khoshnegah *et al.* were the first to report the detection of *H. canis* infection in dogs in Iran. The diagnosis was confirmed under a microscope by identifying *H. canis* gametocytes within neutrophils in peripheral blood smears and bone marrow smears stained with Giemsa (Khoshnegah *et al.*, 2009). Zeinali *et al.* made a significant contribution by documenting the first-ever occurrence of *H. canis* infection in dogs living in northwestern Iran. The results of their research showed a significant occurrence of canine hepatozoonosis in the examined area, suggesting that the disease is widespread among dogs in that particular region (Zeinali *et al.*, 2022). Sanchez *et al.* in 2021 determined the prevalence of Hepatozoon infections in 80 stray dogs in Cuba and confirmed the species identity and phylogenetic relationships of the causative agent. The DNA sequences of the 18S rRNA gene of the *Hepatozoon* spp. from Cuban dogs showed a nucleotide identity >99% with those of 18S rRNA sequences of *H. canis* isolates from Czech Republic, Brazil and Spain. Phylogenetic analysis revealed that obtained sequences clustered within the *H. canis* clade, different from the *Hepatozoon felis* or *H. americanum* clades (Díaz-Sánchez *et al.*, 2021).

### 3. Reports of *Hepatozoon* Species in Foxes

In 2023, Uiterwijk *et al.* conducted a European study which discovered the existence of Hepatozoon in red foxes throughout Europe (Uiterwijk *et al.*, 2023). A study conducted by Cardoso *et al.* in 2014 examined the occurrence of *H. canis* in red foxes from different regions of Portugal.

**Table 1**

The worldwide occurrence of Hepatozoon species in various animal species within the realm of wildlife.

Animal	Number of sampling and animal	Country/ location of sampling	species	The type of material sampled	prevalence	laboratory method	Reference
Dog	274 dogs	USA	Hepatozoon	Blood species	Out of 77 canine blood samples that tested positive for 68 (24.8%) were found to have H. americanum, Hepatozoon spp., 2 (0.7%) had H. canis, Hepatozoon spp., both H. americanum and H. canis.	Nested PCR and real-time PCR	(Allen <i>et al.</i> 2008).
Dog	158 dogs	Brazil	Hepatozoon spp.	Blood samples	The occurrence of H. canis was found to be quite high, at 66.45%.	microscopy and molecular analysis	(de Castro Demoner <i>et al.</i> 2016).
Cat	37 free-roaming domestic cats	Brazil	Anaplasmataceae, Hepatozoon sp. And.	Blood	Hepatozoon spp. 0%	PCR	(André <i>et al.</i> 2014).
Red fox	91 Red foxes (Vulpes vulpes)	Portugal	Hepatozoon canis	Blood and tissues including bone marrow heart, ..	PCR: 68 out of 90 foxes (75.6%)	PCR and histopathological detection	(Cardoso <i>et al.</i> 2022).
Grey wolf	107 grey wolf (Canis lupus L.1758)	Serbia	Hepatozoon canis	Spleen samples	Out of the 30 locations that were tested, 26 of them had confirmed cases of H. canis, with a total of 62 individuals (57.94%) testing positive for the disease.	The 18S ssrRNA gene was analyzed genetically by sequencing a fragment of it using PCR.	(Kuručki, Tomanović, Sukara & Ćirović 2022).
Jackal	An Indian jackal	India	Hepatozoon Canis	Blood sample	After undergoing eight weeks of rehabilitation, the animal experienced a significant decline in its haemoglobin concentration, platelet count, and weight, and was subsequently diagnosed with Hepatozoonosis.	CBC	(Kolangath <i>et al.</i> 2022).
Giant panda	14 giant pandas (Ailuropoda melanoleuca)	USA, china and UK	a novel Hepatozoon species	Blood smear	100% prevalence (14/14) across	-	(Yu <i>et al.</i> 2019).
Short-beaked echidna	18 short-beaked echidna	-	Hepatozoon tachyglossi	Blood samples	Nine (50%) of these animals showed parasitaemia with H tachyglossi.	CBC and multiple biochemical analysis electron microscopy	(Ploeg <i>et al.</i> 2008).
Captive cranes	3 Young captive cranes	Japan	Hepatozoon-like organisms	Liver, spleen, blood films and bone marrow	-	microscope	(YASUDA, KONC & KOYAMA 1987).
Grey squirrel	Grey Squirrel (Sciurus carolinensis Gmelin, 1788)	USA	Hepatozoon griseisciuri n. sp.	-	-	-	(Clark 1958).
chuckwalla Sauromalus spp.	-	USA	Hepatozoon sauromali sp.	Blood	-	Blood film test	(Lewis & Wagner 1964).
Snake	Eighty-three Neotropical Rattlesnake, Crotalus durissus collilineatus	Brazil	Hepatozoon	Blood	Wild snakes that were caught had an infection rate of 38.3%, but no Hepatozoon sp. was found.	spectrophotometry	(Motta <i>et al.</i> 2011).
Snake	1 Southern Water Snake (Nerodia fasciata pictiventris).	USA	Hepatozoon sp.	Peripheral blood	-	Gross and histopathologic examination of the liver	(Wozniak <i>et al.</i> 1998).
wild mammals and ticks	2801 wild mammals and 754 ticks	Europe	Hepatozoon species	Ticks and a piece of spleen From each wild mammal.	Out of the total number of wild mammals observed, 370 (13.2%) were found to be carrying ticks. In the study, a total of 754 ticks were collected and tested for Hepatozoon. The results showed that none of the 35 Hyalomma sp. ticks tested positive for Hepatozoon, while 4 out of 25 Dermacentor spp. ticks (16.0%) and none of the 23 Haemaphysalis spp. ticks tested positive. Additionally, 24 out of 50 Ixodes ticks (5.3%) and 3 out of 221 Rhipicephalus spp. ticks (1.4%) tested positive for Hepatozoon, with an overall positive rate of 4.2% (32/754). The most commonly found species of Hepatozoon was H. canis, with a total of 22 positive cases.	PCR	(Uiterwijk <i>et al.</i> 2023).
Small mammal species	30 Ear samples mammal species	Italy	Hepatozoon spp.	Ear samples	an overall prevalence of 35.9%	PCR	(Ferrari <i>et al.</i> 2022).
Sika deer (Cervus nippon centralis) and Japanese wild boar (Sus scrofa leucomystax)	63 Sika deer (Cervus nippon centralis) and 30 Japanese wild boar (Sus scrofa leucomystax) 63 heads were found.	Japan	Sarcocystis cyst (sarcocyst) and Hepatozoon sp.	-	During the survey, it was discovered that Hepatozoon sp. was present in the muscles of boars.	An examination of the 18S ribosomal DNA through phylogenetic analysis	(Matsuo, Uetsu, Takashima & Abe 2016).

The wild felids and canids found in India include the Asiatic lion, Indian tiger, Indian leopard, Indian wild dog as well as domestic cats that are native to India.	-	India	Hepatozoon spp.	Blood samples	PCR testing revealed the presence of Hepatozoon spp. in the blood samples of several animals. Specifically, out of the 9 Asiatic lions tested, 5 were positive for Hepatozoon spp. Additionally, 2 out of 5 Indian tigers, 2 out of 4 Indian leopards, and 2 out of 2 Indian wild dogs were also found to have the presence of Hepatozoon spp. Furthermore, 2 out of 4 domestic cats and 2 out of 3 domestic dog samples were positive for Hepatozoon spp.	PCR.	(Pawar <i>et al.</i> 2012).
Wild carnivores from Brazil and other exotic locations, including captive wild cat and dogs	In Brazil, there are 165 captive wild felids and 100 captive wild canids, which are re exotic carnivorous animals.	Brazil	Hepatozoon spp.	Blood samples	Six types of cats, including four little spotted cats, one jaguarondi, and one puma, as well as five types of canids, including two bush dogs, one fox, one crab-eating fox, and one maned wolf, were found to be infected with Hepatozoon spp.	PCR.	(André <i>et al.</i> 2010).
36 distinct species of wild animals, including reptiles, birds, and mammals.	A total of 181 wild animals were examined, consisting of 36 distinct species. These species included 2 reptiles, 5 birds, and 29 mammals.	Brazil	Piroplasmida (genera Babesia, Theileria, Cytauxzoon) or Hepatozoon.	Blood samples	The study found various types of Piroplasmida and Hepatozoon agents in different animals. Specifically, Cytauxzoon felis was detected in one ocelot, Theileria cervi in two red brocket deer, Theileria spp. in three nine-banded-armadillos, one agouti, and four lowland pacas, and Babesia spp. in one common opossum and one white-lipped peccary. Additionally, Hepatozoon sp. (possibly Hepatozoon caimani) was found in three spectacled caimans, Hepatozoon felis in an ocelot, and Hepatozoon spp. in one scorpion mud turtle and one lowland paca.	PCR.	(Soares <i>et al.</i> 2017).

The researchers utilized both histopathology and molecular analysis to investigate this prevalence. This research involved the examination of blood and tissue samples obtained from 91 red foxes (*Vulpes vulpes*). PCR and sequencing methods were used to detect *H. canis* in 75.6% of the foxes that were sampled. Histopathological analysis revealed the presence of *H. canis* meronts in the bone marrow and spleen of certain foxes. Ultimately, it was concluded that PCR is more sensitive than histopathological examination in diagnosing *H. canis* (Cardoso *et al.*, 2014). PCR investigations carried out in different countries in southern Europe have shown a considerable prevalence of *H. canis* infection in foxes. In the case of red foxes from northern Spain, the use of blood PCR revealed a significantly lower occurrence of *H. canis* at 28.6% (2 out of 7) compared to the results obtained from this study in Portugal, where it reached 70.5%. When the spleen PCR method was used, the occurrence of *H. canis* in red foxes showed significant increase. In central Spain, the prevalence rate reached as high as 90% (18 out of 20), while in Croatia it was 23.0% (44 out of 191), and in Italy it was 16.4% (16 out of 119) (Gabrielli *et al.*, 2010; Criado-Fornelio *et al.*, 2006; Dežek *et al.*, 2010; Gimenez *et al.*, 2009). During a previous study carried out in southern Portugal, it was noted that among the 301 red foxes analyzed using blood, bone marrow, and spleen and lymph node smears under a microscope, a considerable proportion of 143 (48%) were found to be infected with meronts or gamonts (Conceição *et al.*, 1988). Meanwhile, in Israel, an enzyme-linked immunosorbent assay was employed to test 84 fox sera, and it was found that 20 (23.8%) of them exhibited positive results for antibodies to *H. canis* (Fishman *et al.*, 2004). The differences in research findings from vari-

ous studies might suggest that the rates of occurrence vary depending on the geographical location. Nevertheless, these disparities could also arise due to the usage of different methods, like serology for identifying exposure and PCR for detecting infection, as well as variations in the tissues sampled (Criado-Fornelio *et al.*, 2007). There have been debates surrounding the detection of *H. canis* in striated muscle. This organism has been identified through PCR analysis in the heart and striated muscle tissues of red foxes from Slovakia (Majláthová *et al.*, 2007). Generally, there is a noticeable difference in the occurrence rates of *Hepatozoon* spp. between red foxes and dogs living in the same areas. *Hepatozoon* spp. is more commonly found in red foxes, while dogs have considerably lower rates of infection (Cassini *et al.*, 2009; Conceição-Silva *et al.*, 1988; Gabrielli *et al.*, 2010).

#### 4. Reports of *Hepatozoon* Species in Wolves, Coyotes, Jackals and Hyenas

Research conducted by Mercer *et al.* in the USA in 1988 discovered that twelve coyotes (*Canis latrans*) from the Texas Gulf Coast had been infected with *Hepatozoon* sp. It is crucial to acknowledge that this study had certain limitations. These limitations encompassed a small sample size and a dearth of long-term follow-up data (Mercer *et al.*, 1988). In a more recent study conducted in 2022 by Kuručki *et al.* a thorough examination was performed on spleen samples taken from 107 gray wolves in Serbia. Using PCR, it was determined that 57.94% of the animals examined tested positive for *H. canis*. The presence of *H. canis* was confirmed in 62 out of 107 individuals (57.94%)

across 30 distinct locations. Genetic analysis has revealed the presence of five unique sequence types of *H. canis* in grey wolves, indicating a notable degree of diversity. The dietary habits of grey wolves in Serbia might play a role in fostering this diversity of *H. canis*. To enhance our comprehension further, further research is needed to explore the transmission mechanisms, potential sources of infection, and the impact of this pathogen on carnivores in the wild. This study is particularly noteworthy as it represents the first confirmation of *H. canis* presence in grey wolf populations in Serbia, highlighting the important role of grey wolves in the spread of this pathogen within natural ecosystems (Kuručki *et al.*, 2022). Hepatozoonosis in wild canids in India has been infrequently documented. In a study conducted by Kolangath *et al.* in 2022, a paraplegic Indian jackal with a decubital ulcer was diagnosed with *H. canis* following rehabilitation. The jackal was effectively treated using doxycycline and received supportive care, which led to a full recovery (Kolangath *et al.*, 2022). In a European study conducted by Uiterwijk *et al.* in 2023, it was found that *H. canis* had the highest prevalence among carnivores, specifically gray wolves and golden jackals, as well as in ticks. Out of the ticks tested, 32 ticks (4.2%) were identified as positive for *Hepatozoon*, with *H. canis* being the most frequently detected species. These findings indicate the potential for the pathogen to spill over into domestic animals, including domestic dogs (Uiterwijk *et al.*, 2023). In certain cases, the occurrence of clinical disease and mortality in hyenas has been associated with *Hepatozoon* spp. infection in young coyotes (East *et al.*, 2008; Garrett *et al.*, 2005; Kocan *et al.*, 2000). A thorough examination of the 18S rRNA of *Hepatozoon* isolates from coyotes in the south-central United States revealed 19 distinct sequences comprising 3 major clusters of *Hepatozoon* spp., i.e., 1 most closely related to *H. americanum*, another most closely related to *H. canis*, and the third an intermediate between the 2 groups and it was found the diversity of *Hepatozoon* spp. in coyotes appears greater than previously recognized (Starkey *et al.*, 2013).

## 5. Reports of *Hepatozoon* Species in Domestic and Wild Felids

**Cats.** In 1988, Mercer *et al.* conducted a study in the USA which revealed that three bobcats (*Lynx rufus*) from the Texas Gulf Coast were infected with *Hepatozoon sp* (Mercer *et al.*, 1988). Nevertheless, the study encountered some limitations, including a limited number of participants and the lack of data for long-term monitoring. In a separate study conducted in northern and central Portugal, the molecular evaluation of 320 cats revealed that none of them tested positive for *H. canis* using PCR analysis. However, it was dis-

covered that 15.6% of the animals were PCR-positive for *H. felis* (Vilhena *et al.*, 2013). In a study conducted by André *et al.* in Brazil in 2014, they examined 37 free-roaming domestic cats using the PCR method, but they were unable to identify any *Hepatozoon* spp. Infections (André *et al.*, 2014). Before this study in Brazil, there were a few instances where phylogenetically related species of *Hepatozoon*, similar to *H. canis* and *H. felis*, had been found in cats residing in the state of São Paulo. Although these occurrences were infrequent. (de Bortoli *et al.*, 2011; Rubini *et al.*, 2006). Further research is crucial to enhance our understanding of how cats contribute to the spread of vector-borne diseases. André *et al.* (2010) conducted a study using molecular techniques to identify *Hepatozoon* spp. in different felids (including four little spotted cats, one jaguarondi, and one puma) and several canids (including two bush dogs, one fox, one crab-eating fox, and one maned wolf) in Brazil. This is the first time that *Hepatozoon* spp. has been identified in pumas, jaguarondis, bush dogs, and maned wolves in Brazil. Using PCR analysis, it was found that the prevalence of *Hepatozoon* spp. in wild felids in Brazil is lower compared to that in Neotropical felids in Maranhao and Ceara, domestic cats in Barcelona, and Iriomote wild cats and Tsushima leopard cats in Japan. The study also found that various species or genotypes of *Hepatozoon* spp. could infect wild carnivores in Brazil. The existence of *Hepatozoon* spp. in endangered Brazilian wild cats and dogs is a cause for concern due to the possible risks it poses to these animals. This is especially true for individuals with weakened immune systems or concurrent infections (André *et al.*, 2010).

## 6. Reports of *Hepatozoon* Species in Lions, Tigers and Leopards

In a study carried out by Mercer *et al.* in 1988, it was discovered that *Hepatozoon sp.* had infected six ocelots (*Felis pardalis*) from the Texas Gulf Coast (Mercer *et al.*, 1988). This particular wild species has the capability to act as a reservoir of infection for domestic dogs. Nevertheless, it is crucial to acknowledge that the study had certain limitations, such as a small sample size and a lack of long-term follow-up data. Pawar *et al.* conducted a study in 2012 where they identified the presence of *Hepatozoon* spp. in a diverse range of animals, encompassing Asiatic lions, Indian tigers, Indian leopards, Indian wild dogs, domestic cats, and domestic dogs. The specific species of *Hepatozoon* identified varied depending on the animal. *H. felis* was identified as the detected species in Asiatic lions, Bengal tigers, Indian leopards, and domestic cats. On the other side, *H. canis* was found to be the identified species in Indian wild dogs and domestic dogs. The detection of *H. felis* and *H. canis* in both domestic and wild animals

indicates that these species are not confined to specific hosts and possess the capability to infect a range of felids and canids (Pawar *et al.*, 2012). Soares *et al.* conducted a study in 2017 in the Brazilian Amazon, where they identified the presence of *H. felis* in an ocelot (Soares *et al.*, 2017). It is evident based on morphological and molecular data that *Hepatozoon luiperdjien* and *Hepatozoon ingwen* which infect felines such as leopards are distinct species. These species are also distantly related to *H. felis* (GenBank: AY628681) based on 18SrDNA sequence comparisons, isolated by Criado Fornelio *et al.* from domestic cats from Spain. The *H. felis* isolates from Spain are widely regarded as the representative *H. felis* isolates to be used for comparison (Criado-Fornelio *et al.*, 2006).<sup>55</sup> Thus, based on the phylogenetic relationships and comparisons of *H. felis* and *H. felis*-like species of *Hepatozoon*, the identity of the *Hepatozoon*-infecting large carnivores, currently identified as *H. felis* by Pawar *et al.* is questioned.<sup>73</sup>

## 7. Reports of *Hepatozoon* Species in Pandas,

Sloth Bears and Wild Boar

In a separate study by Yu *et al.* in 2019, which focused on the blood of 14 giant pandas, a previously unknown species of *Hepatozoon* was identified in all of the pandas. Notably, all of these animals showed no symptoms of infection at the time of sampling (Yu *et al.*, 2015). In this study, Ribosomal DNA sequences indicated a previously undescribed *Hepatozoon* species. Phylogenetic and distance analyses of the sequences placed it within its own branch, clustered with Old World species with carnivore (primarily ursid and mustelid) hosts. A significant number of Indian sloth bears have been found to be infected with *Hepatozoon ursi*, which poses a potential threat to their health and survival (Pawar *et al.*, 2011). Additionally, a new species of *Hepatozoon* called *Hepatozoon apri* has been discovered in Japanese wild boar (*Sus scrofa leucomystax*), and it is closely related to the *Hepatozoon* species found in carnivores (Yamamoto *et al.*, 2017). Furthermore, in a study carried out by Matsuo *et al.* in Japan in 2016, *Hepatozoon sp.* was discovered during an examination of pig muscles molecularly, and authors suggested *Hepatozoon* isolates from wild boars might be host-specific (Matsuo *et al.*, 2016).

## 8. Reports of *Hepatozoon* Species in Short-Beaked Echidna and Martens

Infection with *Hepatozoon tachyglossi* in echidnas is primarily accidental, with only one weakened individual displaying clinical symptoms. In a study conducted by Ploeg *et al.* in 2008, which examined the blood of 18 short-beaked echidnas, *H. tachyglossi* was detected in

50% of the cases using various biochemical analyses and electron microscopy. However, in this review, authors do not observe consistent serum biochemical changes associated with parasitemia or monocytosis (Ploeg *et al.*, 2008). In short-beaked echidnas, the presence of *H. tachyglossi* as an intracellular parasite may be an accidental discovery during hematologic examination and is unlikely to serve as a reliable prognostic indicator. In order to find out the presence of the mentioned parasite (Hill *et al.*, 2022). The *Hepatozoon* species found in the hearts of Japanese martens is probably the same species found in pine martens in Scotland (Kubo *et al.*, 2009). Pine martens in Scotland were found to be infected with a newly discovered species of *Hepatozoon*, which is closely related to *H. canis*. This infection led to myocarditis and myositis, but did not result in any noticeable illness (Simpson *et al.*, 2005). In a European study carried out by Uiterwijk *et al.* in 2023, it was observed that *H. martis* was predominantly found in stone martens and pine martens (Uiterwijk *et al.*, 2023). The newly identified species, *Hepatozoon martis*, primarily infects martens and has a strong affinity for muscle tissue. It is also capable of transmission through non-vectorial routes such as transplacental, transmission by predation and carnivorism (Hodžić *et al.*, 2013).

## 9. Reports of *Hepatozoon* Species in Small Mammals

*Hepatozoon perniciosum* was discovered in laboratory (white rats) in Washington, D.C. The schizogonic cycle was identified in the liver, while the sporogonic cycle occurred in the mite (*Echinolaelaps echidninus*), which served as the vector (Clark, 1985). Moreover, a sequence resembling *Hepatozoon ayorgbor* was identified in three spleen samples taken from great gerbils in northwestern China (Ji *et al.*, 2021). Several species from the *Hepatozoon* genus, including *H. lavieri*, *H. erhardovae*, *H. sylvatici*, *H. griseisciuri*, and on occasion *H. muris* (formerly known as *H. perniciosum*), have been identified in (European rodents) (Laakkonen *et al.*, 2001; Al-Quraishy *et al.*, 2021; De Monte, 1971; Karbowski *et al.*, 2005; Pawelczyk *et al.*, 2004; Sebek, 1978). In a 2023 European study conducted by Uiterwijk *et al.*, rodents were found to harbor *Hepatozoon sp.* and *H. ayorgbor* (Uiterwijk *et al.*, 2023). In a study conducted in the North-Eastern Italian Alps, *Hepatozoon* spp. was found in 35.9% of the 830 ear samples collected from 11 various small mammal species. Among these species, *My. glareolus* and *Apodemus* spp. demonstrated the highest rates of infection. Significantly, this study represented the initial identification of *Hepatozoon* spp. in *C. nivalis*, an alpine species (Ferrari *et al.*, 2022). *H. canis*, a parasite transmitted by ticks, is prevalent in rural areas of Brazil. Al-

though the specific vectors responsible for transmitting *H. canis* remain unidentified, this has sparked speculation about alternative methods of transmission. In forest fragments located near rural regions in Botucatu County, Brazil, it was found that small rodents were harboring distinct *Hepatozoon* spp. infections that were different from *H. canis*. While sylvatic rodents may not function as reservoirs for *H. canis*, they have the potential to act as paratenic hosts for *Hepatozoon* spp. This means they can carry infective stages of the parasite, which can subsequently be transmitted to intermediate host predators (Demoner *et al.*, 2016). A high prevalence of *Hepatozoon* infection was observed among wild rodents in southeastern Brazil, reaching 55%. Merino *et al.* conducted a study in 2009 where unidentified species of *Hepatozoon* were detected in 47% of 17 rodents, namely (*Abrothrix olivaceus*) and (*Abrothrix sanborni*) (Merino *et al.*, 2009). Likewise, in 2014, Maia *et al.* reported a prevalence rate of 41% for *Hepatozoon* parasites among wild rodents from North Africa (Maia *et al.*, 2014). *Hepatozoon* spp. are commonly found in wild rodents in North America. For example, a study conducted by Johnson *et al.* in 2007 discovered *Hepatozoon* sp. in 58% of cotton rats and 33.3% of white-footed mice in the United States (Johnson *et al.*, 2007). A groundbreaking study conducted by Wolf *et al.* in 2016 revealed the first-ever identification of *Hepatozoon* sp. in sylvatic rodents in Brazil. The study found that the prevalence of (*Calomys callosus*) rodents was relatively low, specifically at 7% (Wolf *et al.*, 2016). The *Hepatozoon* genus, was initially documented in grey squirrels in Maryland, USA, as well as in California ground squirrels in the nearctic region. Multiple species of *Hepatozoon* have been identified within the Sciuridae family; however, specific differentiation remains unclear due to limited knowledge regarding their life cycles. The detection of *Hepatozoon* in grey squirrels in Maryland presented a challenge due to the coexistence of Miller's species in the same area. Further investigation is necessary to understand the sexual and asexual cycles of *Hepatozoon* within arthropod vectors and infected squirrels. This research will aid in determining the exact identity of *Hepatozoon* found in grey squirrels (Clark, 1958). The role of rodents in the epidemiology of *Hepatozoon* sp. Should be further investigated to evaluate if genotypes circulating in rodents are specific of this group of mammals or might also circulate in wild carnivores and reptiles.

*Hepatozoon* sequences detected in rodents in Slovakia and the Czech Republic were phylogenetically related to sequences detected in lizards and snakes, positioning apart from *H. canis* detected in dogs (Hamšíková *et al.*, 2016). Sousa obtained similar results when analyzing *Hepatozoon* sequences from rodents sampled in Brazilian Pantanal. These results suggest that rodents might play an important role as intermediate or paratenic hosts for *Hepatozoon* infec-

tions in reptiles (de Sousa *et al.*, 2017). The sequences detected in rodents by Parles *et al.* in Brazil in 2019 were positioned in a large clade comprising *Hepatozoon* sequences previously detected in rodents and reptiles. In addition, *Hepatozoon* sequences from felids and canids were grouped in another large clade. They reported a variety of *Hepatozoon* haplotypes associated with small mammals in three Brazilian biomes: Cerrado, Atlantic Forest, and Pantanal. Through phylogenetic analyses, the *Hepatozoon* agents grouped in the rodent-marsupial-reptile large clade of *Hepatozoon* spp. from the world (Perles *et al.*, 2019).

In the latter case, rodents would play a role as intermediate or paratenic hosts in the epidemiological cycles.

## 10. Reports of *Hepatozoon* Species in Wild Bbirds

The identification of *Hepatozoon* species in birds has been discussed in multiple studies (Bennett *et al.*, 1992; Merino *et al.*, 2014). In a study conducted by Shimizu *et al.* in 1987, microscopic examination of the liver, spleen, blood film, and bone marrow of captive cranes in Japan revealed the presence of *Hepatozoan-like* organisms. These organisms were associated with significant clinical symptoms, such as hepatomegaly (enlarged liver) and splenomegaly (enlarged spleen), in these animals (Shimizu *et al.*, 1978). Birds of prey can also become infected with *Hepatozoon* by consuming rodent baits that contain the parasites (Peirce *et al.*, 2005). In Tanzania, a previously unknown species of avian blood parasite called *Hepatozoon ellisgreineri* was recently identified in (secretary birds). What sets this species apart is its unique development within granulocytes, particularly heterophils, which distinguishes it from other *Hepatozoon* parasites found in birds. This discovery significantly enhances our understanding of avian *Hepatozoon* species and other blood parasites in birds, thereby advancing our knowledge of the diverse range of haematozoa (Valkiūnas *et al.*, 2016). In seabirds, blood parasites of the genus *Hepatozoon* had first been found in 4 species of albatrosses (*Hepatozoon albatrossi*). Subsequently, an undetermined *Hepatozoon* sp. was reported from 2 seabird species breeding at offshore islands of northern Mexico: 1 of 15 magnificent frigate birds *Fregata magnificent* and 1 of 5 black storm petrels *Oceanodroma Melania* (Clark *et al.*, 1969). However, neither *H. albatrossi* nor the undetermined *Hepatozoon* from Mexico have been mentioned in the later literature, and no molecular data are available. The sequences detected in rodents by Merino *et al.* in USA in 2014 specified that *Hepatozoon peircei* appears phylogenetically clustered with *Hepatozoon* species infecting rodents, marsupials, reptiles, and amphibians and is distant from a

clade including those infecting carnivores and a group of species infecting lizards. However, the position of this avian Hepatozoon is not resolved.

## 11. Reports of *Hepatozoon* Species in Lizards, Snakes and Crocodiles

Lewis and Wagner carried out a study in the USA in 1964, where they discovered a previously unknown species of *Hepatozoon* called *Hepatozoon sauromali* sp. This parasite was found in the blood of (chuckwalla lizards) belonging to the *Sauromalus* species. The study revealed that gametocytes, the sexual stages of the parasite, develop within the red blood cells, while schizogony (asexual reproduction) occurs in the liver. The mature schizont contains an average of 24 merozoites, which are the resulting daughter cells. The parasite is transmitted by a newly identified mite species called *Hirstiella*, and the transmission takes place through the oral route. Sporogony occurs within the body cavity of the mites, resulting in the formation of large oocysts containing around 12 sporozoites. When mites feed on chuckwallas infected with the parasite, they can transmit the infection to “clean” (Sceloporus lizards). Approximately 30 days later, these previously uninfected lizards exhibit blood cell infections (Lewis and Wagner, 1964). In 1938, Lavier and Callot described *H. burneti* as a novel species discovered in the gecko species known as *Tarentola mauritanica*. These researchers hypothesized that a genus of sand fly called *Phlebotomus* sp. served as the invertebrate host responsible for transmitting the parasite (Lavier and Callot, 1938). In 1962, Mackerras identified a novel species called *H. breinli* in a spotted goanna known as (*Varanus tristis orientalis*). The reproductive phases of this parasite were observed to develop within *Culex fatigans* mosquitoes, indicating that this particular mosquito species serves as the vector for *H. breinli* (Mackerras, 1962). *Hepatozoon* spp. has been detected in 8% of ticks found on mammals and 96% of ticks found on snakes in Thailand (Sumrandee *et al.*, 2015). In Shanghai, China, (King rat snakes) have been found to be infected with a new species of *Hepatozoon* called *Hepatozoon chinensis*. This species is unique from other known *Hepatozoon* species and alters the shape of the erythrocytes that it infects (Han *et al.*, 2015). In a study conducted by Wozniak *et al.* in 1998 in the USA, *Hepatozoon* sp. was identified in the bloodstream of a southern water snake (*Nerodia fasciata pictiventris*). One of the clinical symptoms observed in this case was the presence of granulomatous hepatitis, which was associated with *Hepatozoon* sp (Wozniak *et al.*, 1998). In 2011, Motta *et al.* conducted a study in Brazil, focusing on eighty-three neotropical rattlesnakes known as *Crotalus durissus collilineatus*. The researchers found that the infection rate of *Hepa-*

*tozoon* spp. in wild-caught snakes was 38.3%, whereas no infection was detected in captive-bred snakes. It was observed that infected snakes tended to have lower weight, length, and weight-length ratios. Furthermore, it was observed that these infected snakes had an increase in immature red blood cells (Motta *et al.*, 2011). In 2017, Soares *et al.* conducted a study in the Brazilian Amazon which discovered the existence of *Hepatozoon* sp. in (spectacled caimans).<sup>74</sup> In 2006, a study conducted by Vilcins *et al.* related to blood smears of 5 reptiles, including the water python, and their ticks examined for the presence of the *Hepatozoon* sp. DNA was detected in all tick and reptile species, with 57.7% of tick samples (n = 187) and 35.6% of blood smears (n=35) showing evidence of infection (Vilcins *et al.*, 2009). Tome *et al.* 2016 analyzed nearly 600 samples from several species of the Wall Gecko genus *Tarentola* of the Mediterranean Basin and the Canary Islands for the presence of these parasites using haemogregarine-specific primers to assess the diversity and distribution of *Hepatozoon* spp. in this group of hosts. Their results show that geckos of *Tarentola* spp. harbor a great diversity of haemogregarines (Tomé *et al.*, 2016). Junsiri *et al.* in Thailand identify and evaluate the phylogenetic relationships of *Hepatozoon* sp. in crocodile monitors. The phylogenetic results revealed that *Hepatozoon* sp. in crocodile monitors had 99% similarity with *H. caimani*, and the *Hepatozoon* 18S rRNA gene was grouped in the same clade as crocodiles, reptiles, rodents, and amphibians (Junsiri *et al.*, 2024). In a study conducted by Bouer *et al.* in 2017 in the Brazil, *Hepatozoon* spp.-gametocytes were found in 70.8% (17/24) and 88.8% (8/9) of blood smears from free-ranging and captive caimans (*Caiman crocodilus yacare*), respectively. *Hepatozoon* spp. 18S rRNA DNA was found in 79.2% (19/24) and 88.8% (8/9) of free-ranging and captive caimans, respectively.

## 12. Conclusion and future perspective

Apicomplexa, a group of protozoa, continues to be one of the least explored groups, making accurate identification of *Hepatozoon* spp. challenging. To tackle this issue, we carried out an extensive investigation to establish how common *Hepatozoon* spp. is among different types of animals living in the wild or in captivity in various parts of the globe. This study examined case reports and articles that used PCR-based methods with sequencing to identify *Hepatozoon* spp. The epidemiology of *Hepatozoon* spp. In wild and domestic animals is still not fully elucidated around the world. According to the research, *Hepatozoon* spp. infections are prevalent in alpine environments and are commonly found in various species of small mammals that reside in these regions. The prevalence of *Hepatozoon* spp. varies across different mammal species, with the

bank vole (*My. glareolus*) exhibiting the highest prevalence, followed by the yellow-necked mouse (*A. flavicollis*). Additionally, research has shown that *Hepatozoon* spp. can spread through the consumption of infected vectors, indicating that it serves as a vector-borne pathogen for small mammals. The discovery of *Hepatozoon* spp. in different types of mammals, including those that are domesticated and those that are wild, indicates that there is a possibility of diseases spreading between animals that live in the wild and those that are domesticated. Studies have shown that *H. canis*, a parasite that affects dogs, is present in North America. This discovery contradicts previous beliefs that the parasite was not found in the region. The occurrence of *Hepatozoon* spp. varies among mammal species in different regions, with a higher prevalence observed in Croatia and the Netherlands/Belgium compared to Austria and Bosnia. These studies underscore the significance of comprehending the prevalence and transmission patterns of *Hepatozoon* spp. to evaluate the risk of infection and potential consequences for wildlife and domestic animals. Our research has found that wildlife can be a potential source of infection in environmental settings. Moreover, they can spread parasites to both humans and domestic animals. With the rise in deforestation and urbanization, and the increasing proximity between humans, domestic animals, and wildlife, it is imperative to conduct further research in this field to prevent any outbreaks.

#### Acknowledgments

None

#### Conflicts of interest

The authors affirm that they do not have any known conflicts of interest or personal relationships that could have influenced the work presented in this paper.

#### References

- [1] Alencar NX, Kohayagawa A, Santarém VA. Hepatozoon canis infection of wild carnivores in Brazil. *Veterinary Parasitology*. 1997 Jul 1;70(4):279-82.
- [2] Allen KE, Li Y, Kaltenboeck B, Johnson EM, Reichard MV, Panciera RJ, Little SE. Diversity of Hepatozoon species in naturally infected dogs in the southern United States. *Veterinary Parasitology*. 2008 Jul 4;154(3-4):220-5.
- [3] Al-Quraishy S, Abdel-Ghaffar F, Dkhil MA, Abdel-Gaber R. Haemogregarines and criteria for identification. *Animals*. 2021 Jan 12;11(1):170.
- [4] André MR, Adania CH, Teixeira RH, Vargas GH, Falcade M, Sousa L, Salles AR, Allegretti SM, Felipe PA, Machado RZ. Molecular detection of Hepatozoon spp. in Brazilian and exotic wild carnivores. *Veterinary Parasitology*. 2010 Oct 11;173(1-2):134-8.
- [5] Andre MR, Denardi NC, de Sousa KC, Goncalves LR, Henrique PC, Ontivero CR, Gonzalez IH, Nery CV, Chagas CR, Monticelli C, de Santis AC. Arthropod-borne pathogens circulating in free-roaming domestic cats in a zoo environment in Brazil. *Ticks and tick-borne diseases*. 2014 Sep 1;5(5):545-51.
- [6] Baker HJ, Lindsey JR, Wesibroth SH, editors. *The laboratory rat: biology and diseases*. Elsevier; 2013 Oct 2.
- [7] Baneth GA, Samish M, Alekseev E, Aroch I, Shkap V. Transmission of Hepatozoon canis to dogs by naturally-fed or percutaneously-injected Rhipicephalus sanguineus ticks. *Journal of Parasitology*. 2001 Jun 1;87(3):606-11.
- [8] Battisti E, Zanet S, Khalili S, Trisciuoglio A, Hertel B, Ferroglio E. Molecular survey on vector-borne pathogens in alpine wild carnivores. *Frontiers in Veterinary Science*. 2020 Jan 23;7:1.
- [9] Bennett GF, Earlé RA, Penzhorn BL. Ornithodoros peringueyi (Argasidae) and Xenopsylla trispinis (Siphonaptera), probable intermediate hosts of Hepatozoon atticorae of the South African Cliff Swallow, Hirundo spilodera. *Canadian Journal of Zoology*. 1992 Jan 1;70(1):188-90.
- [10] Brumpt E. Contribution à l'étude d'hepatozoon muris. Utilisation du xénodiagnostic pour l'identification des espèces d'hémogregarines. *Annales de Parasitologie Humaine et Comparée*. 1946;21(1-2):1-24.
- [11] Cardoso L, Cortes HC, Eyal O, Reis A, Lopes AP, Vila-Viçosa MJ, Rodrigues PA, Baneth G. Molecular and histopathological detection of Hepatozoon canis in red foxes (Vulpes vulpes) from Portugal. *Parasites & Vectors*. 2014 Mar 24;7(1):113.
- [12] Cassini R, Zanutto S, Di Regalbono AF, Gabrielli S, Calderini P, Moretti A, Tampieri MP, Pietrobelli M. Canine piroplasmiasis in Italy: epidemiological aspects in vertebrate and invertebrate hosts. *Veterinary parasitology*. 2009 Oct 28;165(1-2):30-5.
- [13] Christophers SS. *The sexual cycle of Leucocytozoon canis in the tick*. Office of the Superintendent of Government Printing, India; 1907.
- [14] Clark GM. Hepatozoon griseisciuri n. sp.; a new species of Hepatozoon from the grey squirrel (Sciurus carolinensis Gmelin, 1788), with studies on the life cycle. *The Journal of parasitology*. 1958 Feb 1;44(1):52-63.

- [15] Clark GW, Swinehart BR. Avian haematozoa from the offshore islands of northern Mexico. *Bulletin of the Wildlife Disease Association*. 1969 Apr 1;5(2):111-2.
- [16] Conceição-Silva FM, Abranches P, Silva-Pereira MC, Janz JG. Hepatozoonosis in foxes from Portugal. *Journal of wildlife diseases*. 1988 Apr;24(2):344-7.
- [17] Craig T, Smallwood J, Knauer K, et al. *H. canis* infection in dogs: clinical, radiographic, and hematologic findings. *J Am Vet Med Assoc* 1978;173(8):967-972.
- [18] Criado-Fornelio A, Rey-Valeiron C, Buling A, Barba-Carretero JC, Jefferies R, Irwin P. New advances in molecular epizootiology of canine hematic protozoa from Venezuela, Thailand and Spain. *Veterinary parasitology*. 2007 Mar 31;144(3-4):261-9.
- [19] Criado-Fornelio A, Ruas JL, Casado N, Farias NA, Soares MP, Müller G, Brum JG, Berne ME, Buling-Saraña A, Barba-Carretero JC. New molecular data on mammalian Hepatozoon species (Apicomplexa: Adeleorina) from Brazil and Spain. *Journal of Parasitology*. 2006 Feb 1;92(1):93-9.
- [20] Cunningham MW, Yabsley MJ. *Primer on tick-borne diseases in exotic carnivores*. Fowlers Zoo and Wild Animal Medicine Current Therapy. Philadelphia, USA, Saunders. 2011 Jul 11;7:458-64.
- [21] de Bortoli CP, André MR, Braga MD, Machado RZ. Molecular characterization of Hepatozoon sp. in cats from São Luís Island, Maranhão, northeastern Brazil. *Parasitology Research*. 2011 Oct;109(4):1189-92.
- [22] de Castro Demoner L, Magro NM, da Silva MR, de Paula Antunes JM, Calabuig CI, ODwyer LH. Hepatozoon spp. infections in wild rodents in an area of endemic canine hepatozoonosis in southeastern Brazil. *Ticks and tick-borne diseases*. 2016 Jul 1;7(5):859-64.
- [23] de Castro Demoner L, Magro NM, da Silva MR, de Paula Antunes JM, Calabuig CI, ODwyer LH. Hepatozoon spp. infections in wild rodents in an area of endemic canine hepatozoonosis in southeastern Brazil. *Ticks and tick-borne diseases*. 2016 Jul 1;7(5):859-64.
- [24] De Monte T. Infection with *Hepatozoon muris* Balf. (Protozoa, Adeleidae) in wild varieties of *Rattus norvegicus* in Trieste. *Schweizer Archiv für Tierheilkunde*. 1971 Nov 1;113(11):654-9.
- [25] de Sousa KC, Fernandes MP, Herrera HM, Ben-evenute JL, Santos FM, Rocha FL, Barreto WT, Macedo GC, Campos JB, Martins TF, de Andrade Pinto PC. Molecular detection of Hepatozoon spp. in domestic dogs and wild mammals in southern Pantanal, Brazil with implications in the transmission route. *Veterinary Parasitology*. 2017 Apr 15;237:37-46.
- [26] Dežek D, Vojta L, Ćurković S, Lipej Z, Mihaljević Ž, Cvetnić Ž, Beck R. Molecular detection of *Theileria annae* and *Hepatozoon canis* in foxes (*Vulpes vulpes*) in Croatia. *Veterinary parasitology*. 2010 Sep 20;172(3-4):333-6.
- [27] Diakou A, Dimzas D, Astaras C, Savvas I, Di Cesare A, Morelli S, Neofitos , Migli D, Traversa D. Clinical investigations and treatment outcome in a European wildcat (*Felis silvestris silvestris*) infected by cardio-pulmonary nematodes. *Veterinary Parasitology: Regional Studies and Reports*. 2020 Jan 1; 19: 100357.
- [28] Díaz-Sánchez AA, Hofmann-Lehmann R, Meli ML, Roblejo-Arias L, Fonseca-Rodríguez O, Castillo AP, Cañizares EV, Rivero EL, Chilton NB, Corona-González B. Molecular detection and characterization of *Hepatozoon canis* in stray dogs from Cuba. *Parasitology international*. 2021 Feb 1;80:102200.
- [29] Duscher GG, Fuehrer HP, Kübber-Heiss A. Fox on the run—molecular surveillance of fox blood and tissue for the occurrence of tick-borne pathogens in Austria. *Parasites & vectors*. 2014 Nov 21;7(1):521.
- [30] Duscher GG, Leschnik M, Fuehrer HP, Joachim A. Wildlife reservoirs for vector-borne canine, feline and zoonotic infections in Austria. *International Journal for Parasitology: Parasites and Wildlife*. 2015 Apr 1;4(1):88-96.
- [31] East ML, Wibbelt G, Lieckfeldt D, Ludwig A, Goller K, Wilhelm K, Schares G, Thierer D, Hofer H. A Hepatozoon species genetically distinct from *H. canis* infecting spotted hyenas in the Serengeti ecosystem, Tanzania. *Journal of Wildlife Diseases*. 2008 Jan 1;44(1):45-52.
- [32] Ezeokoli CD, Ogunkoya AB, Abdullahi R, Tekdek LB, Sannusi A, Ilemobade AA. Clinical and epidemiological studies on canine hepatozoonosis in Zaria, Nigeria. *Journal of Small Animal Practice*. 1983 Jul;24(7):455-60.
- [33] Ferrari G, Girardi M, Cagnacci F, Devineau O, Tagliapietra V. First record of *Hepatozoon* spp. in alpine wild rodents: Implications and perspectives for transmission dynamics across the food web. *Microorganisms*. 2022 Mar 25;10(4):712.

- [34] Fishman Z, Gonen L, Harrus S, Strauss-Ayali D, King R, Baneth G. A serosurvey of Hepatozoon canis and Ehrlichia canis antibodies in wild red foxes (*Vulpes vulpes*) from Israel. *Veterinary Parasitology*. 2004 Jan 5;119(1):21-6.
- [35] Forlano MD, Teixeira KR, Scofield A, Elisei C, Yotoko KS, Fernandes KR, Linhares GF, Ewing SA, Massard CL. Molecular characterization of Hepatozoon sp. from Brazilian dogs and its phylogenetic relationship with other Hepatozoon spp. *Veterinary Parasitology*. 2007 Apr 10;145(1-2):21-30.
- [36] Gabrielli S, Kumlien S, Calderini P, Brozzi A, Iori A, Cancrini G. The first report of Hepatozoon canis identified in *Vulpes vulpes* and ticks from Italy. *Vector-borne and Zoonotic Diseases*. 2010 Nov 1;10(9):855-9.
- [37] Garcia-Vozmediano A, Krawczyk AI, Sprong H, Rossi L, Ramassa E, Tomassone L. Ticks climb the mountains: Ixodid tick infestation and infection by tick-borne pathogens in the Western Alps. *Ticks and Tick-borne Diseases*. 2020 Sep 1;11(5):101489.
- [38] Garrett JJ, Kocan AA, Reichard MV, Panciera RJ, Bahr RJ, Ewing SA. Experimental infection of adult and juvenile coyotes with domestic dog and wild coyote isolates of Hepatozoon americanum (Apicomplexa: Adeleorina). *Journal of Wildlife Diseases*. 2005 Jul 1;41(3):588-92.
- [39] Gevrey J. Hepatozoonose canine. *Rec Med Vet* 1993;169(1):6.
- [40] Gilman SE, Urban MC, Tewksbury J, Gilchrist GW, Holt RD. A framework for community interactions under climate change. *Trends in ecology & evolution*. 2010 Jun 1;25(6):325-31.
- [41] Gimenez C, Casado N, Criado-Fornelio Á, de Miguel FÁ, Dominguez-Peñafiel G. A molecular survey of Piroplasmida and Hepatozoon isolated from domestic and wild animals in Burgos (northern Spain). *Veterinary Parasitology*. 2009 May 26;162(1-2):147-50.
- [42] Hamšíková Z, Silaghi C, Rudolf I, Venclíková K, Mahríková L, Slovák M, Mendel J, Blažejová H, Berthová L, Kocianová E, Hubálek Z. Molecular detection and phylogenetic analysis of Hepatozoon spp. in questing Ixodes ricinus ticks and rodents from Slovakia and Czech Republic. *Parasitology research*. 2016 Oct;115(10):3897-904.
- [43] Han H, Wu Y, Dong H, Zhu S, Li L, Zhao Q, Wu D, Pei E, Wang Y, Huang B. First report of Hepatozoon (Apicomplexa: Adeleorina) from king ratsnakes (*Elaphe carinata*) in Shanghai, with description of a new species. *Acta Parasitologica*. 2015 Apr;60(2):266-74.
- [44] Hill AG, Boyd SP, Clark NJ, Proboste T. Haemoprotozoa in wild shortbeaked echidnas (*Tachyglossus aculeatus*). *Australian Veterinary Journal*. 2022 Mar;100(3):130-4.
- [45] HHodžić A, Alić A, Beck R, Beck A, Huber D, Otranto D, Baneth G, Duscher GG. Hepatozoon martis n. sp. (Adeleorina: Hepatozoidae): Morphological and pathological features of a Hepatozoon species infecting martens (family Mustelidae). *Ticks and tick-borne diseases*. 2018 May 1;9(4):912-20.
- [46] Hoogstraal H. The life cycle and incidence of Hepatozoon balfouri (Laveran, 1905) in Egyptian jerboas (*Jaculus* spp.) and mites (*Haemolaelaps aegyptius* Keegan, 1956). *The Journal of Protozoology*. 1961 Aug;8(3):231-48.
- [47] Inokuma H, Okuda M, Ohno K, Shimoda K, Onishi T. Analysis of the 18S rRNA gene sequence of a Hepatozoon detected in two Japanese dogs. *Veterinary Parasitology*. 2002 Jun 26;106(3):265-71.
- [48] Ivanov A, Kanakov D. First case of canine hepatozoonosis in Bulgaria. *Bulg J Vet Med*. 2003 May 28;6:43-6.
- [49] James SP. On a parasite found in the white corpuscles of the blood of dogs. Office of the Superintendent of Government Printing, India; 1905.
- [50] Ji N, Chen X, Liu G, Zhao S, Tan W, Liu G, Zhang J, Wang Y. Theileria, Hepatozoon and Taenia infection in great gerbils (*Rhombomys opimus*) in northwestern China. *International Journal for Parasitology: Parasites and Wildlife*. 2021 Aug 1;15:79-86.
- [51] Jittapalapong S, Rungphisutthipongse O, Maruyama S, Schaefer JJ, Stich RW. Detection of Hepatozoon canis in stray dogs and cats in Bangkok, Thailand. *Annals of the New York Academy of Sciences*. 2006 Oct;1081(1):479-88.
- [52] Johnson EM, Allen KE, Breshears MA, Panciera RJ, Little SE, Ewing SA. Experimental transmission of Hepatozoon americanum to rodents. *Veterinary parasitology*. 2008 Feb 14;151(2-4):164-9.
- [53] Johnson EM, Allen KE, Panciera RJ, Ewing SA, Little SE, Reichard MV. Field survey of rodents for Hepatozoon infections in an endemic focus of American canine hepatozoonosis. *Veterinary Parasitology*. 2007 Nov 30;150(1-2):27-32.

- [54] Johnson PT, Dobson A, Lafferty KD, Marcogliese DJ, Memmott J, Orlofske SA, Poulin R, Thieltges DW. When parasites become prey: ecological and epidemiological significance of eating parasites. *Trends in ecology & evolution*. 2010 Jun 1;25(6):362-71.
- [55] Junsiri W, Kamkong P, Taweethavonsawat P. First molecular detection and genetic diversity of Hepatozoon sp. (Apicomplexa) and Brugia sp. (Nematoda) in a crocodile monitor in Nakhon Pathom, Thailand. *Scientific reports*. 2024 Feb 12;14(1):3526.
- [56] Karbowski G, Rychlik L, Nowakowski W, Wita I. Natural infections of small mammals with blood parasites on the borderland of boreal and temperate forest zones. *Acta Theriologica*. 2005 Mar;50(1):31-42.
- [57] Khoshnegah J, Mohri M, Movassaghi AR, Mehrjerdi HK. The first report of Hepatozoon canis infection of a dog in Iran. *Comparative Clinical Pathology*. 2009 Nov;18(4):455-8.
- [58] Kocan AA, Cummings CA, Panciera RJ, Mathew JS, Ewing SA, Barker RW. Naturally occurring and experimentally transmitted Hepatozoon americanum in coyotes from Oklahoma. *Journal of Wildlife Diseases*. 2000 Jan 1;36(1):149-53.
- [59] Kolangath SM, Upadhye SV, Dhoot VM, Pawshe MD, Shalini AS, Kolangath RM. Molecular investigation and clinical management of Hepatozoon Canis infection in an Indian jackala case report. *BMC Veterinary Research*. 2022 Apr 20;18(1):144.
- [60] Kubo M, Nagataki M, Agatsuma T, Sakai H, Masegi T, Panciera RJ, Yanai T. Parasitological and molecular features of the Hepatozoon species in the myocardium of Japanese Martens (*Martes melampus melampus*). *Journal of Parasitology*. 2009 Dec 1;95(6):1496-502.
- [61] Kuručki M, Tomanović S, Sukara R, Ćirović D. High prevalence and genetic variability of Hepatozoon canis in grey wolf (*Canis lupus* L. 1758) population in Serbia. *Animals*. 2022 Nov 29;12(23):3335.
- [62] Laakkonen J, Sukura A, Oksanen A, Henttonen H, Soveri T. Haemogregarines of the genus Hepatozoon (Apicomplexa: Adeleina) in rodents from northern Europe. *Folia parasitologica*. 2001 Jan 1;48(4):263-7.
- [63] Lavier G, Callot J. Hepatozoon burnandi n. sp, hemogregarine parasite de Tarentola mauritanica. *Archives de l'Institut Pasteur de Tunis*. 1938;27:444-8.
- [64] Lewis JE, Wagner ED. Hepatozoon sauromali sp. n., a hemogregarine from the chuckwalla (*Sauromalus* spp.) with notes on the life history. *The Journal of Parasitology*. 1964 Feb 1:11-4.
- [65] Lindblad-Toh K, Wade CM, Mikkelsen TS, Karlsson EK, Jaffe DB, Kamal M, Clamp M, Chang JL, Kulbokas III EJ, Zody MC, Mauceli E. Genome sequence, comparative analysis and haplotype structure of the domestic dog. *Nature*. 2005 Dec 8;438(7069):803-19.
- [66] Mackerras MJ. The life of a Hepatozoon (Sporozoa: Adeleidea) of varanid Lizards in Australia. *Australian Journal of Zoology*. 1962 Mar 1;10(1):35-44.
- [67] Maia JP, Álvares F, Boratyński Z, Brito JC, Leite JV, Harris DJ. Molecular assessment of Hepatozoon (Apicomplexa: Adeleorina) infections in wild canids and rodents from North Africa, with implications for transmission dynamics across taxonomic groups. *Journal of Wildlife Diseases*. 2014 Oct 1;50(4):837-48.
- [68] Majláthová V, Hurníková Z, Majláth I, Peško B. Hepatozoon canis infection in Slovakia: imported or autochthonous?. *Vector-Borne and Zoonotic Diseases*. 2007 Jun 1;7(2):199-202.
- [69] Matsuo K, Uetsu H, Takashima Y, Abe N. High occurrence of Sarcocystis infection in Sika deer *Cervus nippon centralis* and Japanese wild boar *Sus scrofa leucomystax* and molecular characterization of Sarcocystis and Hepatozoon isolates from their muscles.
- [70] Menn B, Lorentz S, Naucke TJ. Imported and travelling dogs as carriers of canine vector-borne pathogens in Germany. *Parasites vectors*. 2010 Apr 8;3(1):34.
- [71] Mercer SH, Jones LP, Rappole JH, Twedt D, Laack LL, Craig TM. Hepatozoon sp. in wild carnivores in Texas. *Journal of Wildlife Diseases*. 1988 Jul;24(3):574-6.
- [72] Merino S, Martínez J, Masello JF, Bedolla Y, Quillfeldt P. First molecular characterization of a Hepatozoon species (Apicomplexa: Hepatozoidae) infecting birds and description of a new species infecting storm petrels (Aves: Hydrobatidae). *The Journal of parasitology*. 2014 Jun 1;100(3):338-43.
- [73] Merino S, Vásquez RA, Martínez J, Celis-Diez JL, Gutiérrez-Jiménez L, Ippi S, Sánchez-Monsalvez I, Martínez-De La Puente J. Molecular characterization of an ancient Hepatozoon species parasitizing the 'living fossil' marsupial 'Monito del

- Monte'Dromiciops gliroides from Chile. Biological Journal of the Linnean Society. 2009 Nov 1;98(3):568-76.
- [74] Miller WW. Hepatozoon perniciosum (ng, n. sp.); a haemogregarine pathogenic for white rats; with a description of the sexual cycle in the intermediate host, a mite (Lelaps echidninus). (No Title). 1908.
- [75] Modrý D, Hofmannová L, Papežík P, Majerová K, Votýpka J, Hönig V, Růžek D, Hrazdilová K. Hepatozoon in Eurasian red squirrels *Sciurus vulgaris*, its taxonomic identity, and phylogenetic placement. Parasitology Research. 2021 Aug;120(8):2989-93.
- [76] Motta RO, Cunha LM, Leite RC, Da Silva IJ, Pinto AC, Braga M, Da Cunha AP, De Oliveira PR. Hepatozoon spp. (Apicomplexa: Hepatozoidae) infection and selected hematological values of the neotropical rattlesnake, *Crotalus durissus collilineatus* (Linnaeus, 1758) (Serpentes: Viperidae), from Brazil. Journal of Zoo and Wildlife Medicine. 2011 Sep;42(3):399-407.
- [77] Murata T, Inoue M, Tateyama S, Taura Y, Nakama S. Vertical transmission of Hepatozoon canis in dogs. Journal of Veterinary Medical Science. 1993 Oct 15;55(5):867-8.
- [78] Odwyer LH, Massard CL, de Souza JC. Hepatozoon canis infection associated with dog ticks of rural areas of Rio de Janeiro State, Brazil. Veterinary Parasitology. 2001 Jan 1;94(3):143-50.
- [79] O'Dwyer LH. Brazilian canine hepatozoonosis. Revista brasileira de parasitologia veterinaria. 2011;20:181-93.
- [80] Paperna I, Lainson R. Hepatozoon cf. terzii (Sambon Seligman, 1907) infection in the snake *Boa constrictor constrictor* from north Brazil: transmission to the mosquito *Culex quinquefasciatus* and the lizard *Tropidurus torquatus*. Parasite. 2004 Jun 1;11(2):175-81.
- [81] Pawar RM, Poornachandar A, Arun AS, Manikandan S, Shivaji S. Molecular prevalence and characterization of Hepatozoon ursi infection in Indian sloth bears (*Melursus ursinus*). Veterinary Parasitology. 2011 Dec 15;182(2-4):329-32.
- [82] Pawar RM, Poornachandar A, Srinivas P, Rao KR, Lakshmikantan U, Shivaji S. Molecular characterization of Hepatozoon spp. infection in endangered Indian wild felids and canids. Veterinary Parasitology. 2012 May 25;186(3-4):475-9.
- [83] Pawelczyk A, Bajer A, Behnke JM, Gilbert FS, Sinski E. Factors affecting the component community structure of haemoparasites in common voles (*Microtus arvalis*) from the Mazury Lake District region of Poland. Parasitology Research. 2004 Mar;92(4):270-84.
- [84] Peirce MA. A checklist of the valid avian species of Babesia (Apicomplexa: Piroplasmorida), Haemoproteus, Leucocytozoon (Apicomplexa: Haemosporida), and Hepatozoon (Apicomplexa: Haemogregarinidae). Journal of Natural History. 2005 Dec 23;39(42):3621-32.
- [85] Perles L, Roque AL, DAndrea PS, Lemos ER, Santos AF, Morales AC, Machado RZ, Andr MR. Genetic diversity of Hepatozoon spp. in rodents from Brazil. Scientific Reports. 2019 Jul 12;9(1):10122.
- [86] Ploeg RJ, Nicolson V, Pyne M, Webb R, O'donoghue P. Hepatozoon tachyglossi in the short-beaked echidna (*Tachyglossus aculeatus*). Australian veterinary journal. 2008 May 1;86(5).
- [87] Rajamanickam C, Wiesenhuber E, Zin FM, Hamid J. The incidence of canine haematozoa in Peninsular Malaysia. Veterinary parasitology. 1985 Jan 1;17(2):151-7.
- [88] Rigó K, Majoros G, Szekeres S, Molnár I, Jablonszky M, Majláthová V, Majláth I, Földvári G. Identification of Hepatozoon erhardovae Krampitz, 1964 from bank voles (*Myodes glareolus*) and fleas in Southern Hungary. Parasitology research. 2016 Jun;115(6):2409-13.
- [89] Rizzoli A, Tagliapietra V, Cagnacci F, Marini G, Arnoldi D, Rosso F, Rosà R. Parasites and wildlife in a changing world: The vector-host-pathogen interaction as a learning case. International Journal for Parasitology: Parasites and Wildlife. 2019 Aug 1;9:394-401.
- [90] Rubini AS, dos Santos Paduan K, Perez RR, Ribolla PE, ODwyer LH. Molecular characterization of feline Hepatozoon species from Brazil. Veterinary Parasitology. 2006 Apr 15;137(1-2):168-71.
- [91] Sebek Z. Blood parasites of small mammals in western Hungary. Parasitologia Hungarica. 1978;11:17-22.
- [92] Shimizu T, Yasuda N, Kono I, Koyama T. Fatal infection of Hepatozoon-like organisms in the young captive cranes (*Grus monacha*). Memoirs of the Faculty of Agriculture-Kagoshima University (Japan). 1987;23.
- [93] Shkap V, Baneth G, Pipano E. Circulating antibodies to Hepatozoon canis demonstrated by immunofluorescence. Journal of Veterinary Diagnostic Investigation. 1994 Jan;6(1):121-3.

- [94] Simpson VR, Birtles RJ, Bown KJ, Panciera RJ, Butler H, Davison N. Hepatozoon species infection in wild red squirrels (*Sciurus vulgaris*) on the Isle of Wight. *Veterinary Record*. 2006 Aug;159(7):202-5.
- [95] Simpson VR, Panciera RJ, Hargreaves J, McGarry JW, Scholes SF, Bown KJ, Birtles RJ. Myocarditis and myositis due to infection with Hepatozoon species in pine martens (*Martes martes*) in Scotland. *Veterinary Record*. 2005 Apr;156(14):442-6.
- [96] Soares HS, Marcili A, Barbieri AR, Minervino AH, Moreira TR, Gennari SM, Labruna MB. Novel piroplasmid and Hepatozoon organisms infecting the wildlife of two regions of the Brazilian Amazon. *International Journal for Parasitology: Parasites and Wildlife*. 2017 Aug 1;6(2):115-21.
- [97] Starkey LA, Panciera RJ, Paras K, Allen KE, Reiskind MH, Reichard MV, Johnson EM, Little SE. Genetic diversity of Hepatozoon spp. in coyotes from the south-central United States. *The Journal of Parasitology*. 2013 Apr 1;99(2):375-8.
- [98] Sumrandee C, Baimai V, Trinachartvanit W, Ahantarig A. Hepatozoon and Theileria species detected in ticks collected from mammals and snakes in Thailand. *Ticks and tick-borne diseases*. 2015 Apr 1;6(3):309-15.
- [99] Tomé B, Maia JP, Salvi D, Brito JC, Carretero MA, Perera A, Meimberg H, Harris DJ. Patterns of genetic diversity in Hepatozoon spp. infecting snakes from North Africa and the Mediterranean Basin. *Systematic Parasitology*. 2014 Mar;87(3):249-58.
- [100] Tomé B, Rato C, Harris DJ, Perera A. High diversity of Hepatozoon spp. in geckos of the genus *Tarentola*. *The Journal of Parasitology*. 2016 Aug 1;102(4):476-80.
- [101] Uiterwijk M, Vojta L, Šprem N, Beck A, Jurković D, Kik M, Duscher GG, Hodžić A, Reljić S, Sprong H, Beck R. Diversity of Hepatozoon species in wild mammals and ticks in Europe. *Parasites & Vectors*. 2023 Jan 24;16(1):27.
- [102] Urban MC, Bocedi G, Hendry AP, Mihoub JB, Peer G, Singer A, Bridle JR, Crozier LG, De Meester L, Godsoe W, Gonzalez A. Improving the forecast for biodiversity under climate change. *Science*. 2016 Sep 9;353(6304): aad8466.
- [103] Valkiūnas G, Mobley K, Iezhova TA. Hepatozoon *ellisgreineri* n. sp. (Hepatozoidae): description of the first avian apicomplexan blood parasite inhabiting granulocytes. *Parasitology research*. 2016 Feb;115(2):609-13.
- [104] Vilcins IM, Ujvari B, Old JM, Deane E. Molecular and morphological description of a Hepatozoon species in reptiles and their ticks in the Northern Territory, Australia. *Journal of Parasitology*. 2009 Apr 1;95(2):434-42.
- [105] Vilhena H, Martinez-Díaz VL, Cardoso L, Vieira L, Altet L, Francino O, Pastor J, Silvestre-Ferreira AC. Feline vector-borne pathogens in the north and centre of Portugal. *Parasites & Vectors*. 2013 Apr 15;6(1):99.
- [106] Vincent-Johnson NA, Macintire DK, Lindsay DS, Lenz SD, Baneth G, Shkap V, Blagburn BL. A new Hepatozoon species from dogs: description of the causative agent of canine hepatozoonosis in North America. *The Journal of parasitology*. 1997 Dec 1;1165-72.
- [107] Wolf RW, Aragona M, Munoz-Leal S, Pinto LB, Melo AL, Braga IA, dos Santos Costa J, Martins TF, Marcili A, de Campos Pacheco R, Labruna MB. Novel Babesia and Hepatozoon agents infecting non-volant small mammals in the Brazilian Pantanal, with the first record of the tick *Ornithodoros guaporensis* in Brazil. *Ticks and tick-borne Diseases*. 2016 Apr 1;7(3):449-56.
- [108] Wozniak EJ, Telford Jr SR, DeNardo DF, McLaughlin GL, Butler JF. Granulomatous hepatitis associated with Hepatozoon sp. meronts in a southern water snake (*Nerodia fasciata pictiventris*). *Journal of Zoo and Wildlife Medicine*. 1998 Mar 1:68-71.
- [109] Yamamoto M, Tokiwa T, Tobiume M, Akamatsu S, Matsuo K, Moribe J, Ike K. Hepatozoon *apri* n. sp. (Adeleorina: Hepatozoidae) from the Japanese wild boar *Sus scrofa leucomystax* (Mammalia: Cetartiodactyla). *International Journal for Parasitology: Parasites and Wildlife*. 2017 Dec 1;6(3):354-60.
- [110] Yu JH, Durrant KL, Liu S, Carlin EP, Wang C, Rodriguez J, Bratthauer A, Walsh T, Valitutto MT, Fine L, Murray S. First report of a novel Hepatozoon sp. in giant pandas (*Ailuropoda melanoleuca*). *EcoHealth*. 2019 Jun 15;16(2):338-45.
- [111] Zechmeisterová K, Javanbakht H, Kvičerová J, Široký P. Against growing synonymy: Identification pitfalls of Hepatozoon and Schellackia demonstrated on North Iranian reptiles. *European Journal of Protistology*. 2021 Jun 1;79:125780.
- [112] Zeinali M, Malekifard F, Rakhshanpour A, Yakhchali M. Study on Hepatozoon *canis* in dogs in Urmia region. *Iranian Journal of Animal Science*. 2022 Jan 21;52(4):253-60.